

PRODUCT DATA SHEET

Adipophilin Human, Rabbit Polyclonal Antibody

Cat. No.: RD181161100

Size: 0.1 mg

Source of Antigen: E. coli

Type: Polyclonal Antibody

Host: Rabbit

Other Names:

Perilipin-2, Adipose differentiation-related protein, ADRP, PLIN2, ADFP

Preparation:

The antibody was raised in rabbits by immunization with the recombinant Human Adipophilin.

Amino Acid Sequence of Immunogen:

The immunization antigen (49.03 kDa) is a protein containing 444 AA of recombinant Human Adipophilin. N-Terminal His-tag, 8 extra AA.

MKHHHHHHAS VAVDPQPSVV TRVVNLPLVS STYDLMSSAY LSTKDQYPYL
KSVCEMAENG VKTITSVAMT SALPIIQKLE PQIAVANTYA CKGLDRIER LPILNQPSTQ
IVANAKGAVT GAKDAVTTTV TGAKDSVAST ITGVMDKTKG AVTGSVEKTK SVVSGSINTV
LGSRRMQLVS SGVENALTKS ELLVEQYLPL TEEELEKEAK KVEGFDLVQK PSYYVRLGSL
STKLHSRAYQ QALSRVKEAK QKSQQTISQL HSTVHLIEFA RKNVYSANQK IQDAQDKLYL
SWVEWKRSIG YDDTDESHCA EHIESRTLAI ARNLTQQLQT TCHTLLSNIQ GVPQNIQDQA
KHMGMAGDI YSVFRNAASF KEVSDSLLTS SKGQLQKMKE SLDDVMDYLV
NNTPLNLWLVG PFYPQLTESQ NAQDQGAEMD KSSQETQRSE HKTH

Purification Method:

Immunoaffinity chromatography on a column with immobilized recombinant Human Adipophilin.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard).

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. ****AZIDE FREE****.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C . Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C . Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C . Quality Control Test

Quality Control:

Indirect ELISA – to determine titer of the antibody

SDS PAGE – to determine purity of the antibody

BCA - to determine quantity of the antibody

Applications:

ELISA, Immunohistochemistry, Western blotting

Note:

This product is for research use only.