

## PRODUCT DATA SHEET

### Apoptosis Inhibitor of Macrophage Human, Sheep Polyclonal Antibody

**Cat. No.:** RD184328100

**Size:** 0.1 mg

**Source of Antigen:** E. coli

**Type:** Polyclonal Antibody

**Host:** Sheep

**Other Names:**

CD5 molucel-like protein

**Preparation:**

The antibody was raised in sheep by immunization with the recombinant Human AIM.

**Amino Acid Sequence of Immunogen:**

Total 337 AA. MW: 37.2 kDa (calculated). UniProtKB acc.no. O43866 (Ser20-Gly347).  
N-terminal His-tag (9 extra AA).

MKHHHHHHAS PSGVRLVGGL HRCEGRVEVE QKGQWGTVCD DGWDIKDVAV  
LCRELGCGAA SGTPSGILYE PPAEKEQKVL IQSVSCTGTE DTLAQCEQEE VYDCSHDEDA  
GASCENPESS FSPVPEGVRL ADGPGHCKGR VEVKHQNQWY TVCQTGWSLR  
AAKVVCRLG CGRAVLTQKR CNKHAYGRKP IWLSQMSCSG REATLQDCPS  
GPWGKNTCNH DEDTWVECED PFDLRLVGGD NLCSGRLEVL HkgVWGSVCD  
DNWGEKEDQV VCKQLGCGKS LSPSFRDRKC YGPGVGRWL DNVRCSGEEQ  
SLEQCQHRFW GFHDCTHQED VAVICSG

**Purification Method:**

Immunoaffinity chromatography on a column with immobilized recombinant Human AIM.

**Species Reactivity:**

Human. Not yet tested in other species.

**Antibody Content:**

0.1 mg (determined by BCA method, BSA was used as a standard)

**Formulation:**

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. **\*\*AZIDE FREE\*\***.

**Reconstitution:**

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

**Shipping:**

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

**Storage/Stability:**

The lyophilized antibody remains stable and fully active until the expiry date when stored at  $-20^{\circ}\text{C}$ . Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at  $-80^{\circ}\text{C}$ . Reconstituted antibody can be stored at  $4^{\circ}\text{C}$  for a limited period of time; it does not show decline in activity after one week at  $4^{\circ}\text{C}$ .

**Quality Control:**

Indirect ELISA – to determine titer of the antibody

SDS PAGE – to determine purity of the antibody

BCA - to determine quantity of the antibody

**Applications:**

ELISA, Immunohistochemistry, Western blotting

**Note:**

This product is for research use only.