

PRODUCT DATA SHEET

Procathepsin K Human, Rabbit Polyclonal Antibody

Cat. No.: RD181074100

Size: 0.1 mg

Source of Antigen: E. coli

Type: Polyclonal Antibody

Host: Rabbit

Other Names:

Preparation:

The antibody was raised in rabbits by immunization with the recombinant Human Procathepsin K.

Amino Acid Sequence of Immunogen:

The immunization antigen (36.7 kDa) is a protein containing 314 amino acid residues of recombinant Human Procathepsin K and 12 additional amino acid residues – HisTag (highlighted). The recombinant Human Procathepsin K is without signal peptid (15 amino acid residues).

MRGSHHHHHH GSLYPEEILD THWELWKKTH RKQYNNKVDE ISRRLIWEKN LKYISIHNL
ASLGVHTYEL AMNHLGDMTS EEVVQKMTGL KVPLSHSRSN DTLYIPEWEG
RAPDSVDYRK KGYVTPVKNQ GQCGSCWAFS SVGALEGQLK KKTGKLLNLS
PQNLVDCVSE NDGCGGGYMT NAFQYVQKNR GIDSEDAYPY VGQEESCMYN
PTGKAAKCRG YREIPEGNEK ALKRAVARVG PVSVAIDASL TSFQFYSGV YYDESCNSDN
LNHAVLAVGY GIQKGNKHWI IKNSWGENWG NKG YILMARN KNNACGIANL ASFPKM

Purification Method:

Immunoaffinity chromatography on a column with immobilized recombinant Human Procathepsin K.

Species Reactivity:

Human. Not yet tested in other species.

Antibody Content:

0.1 mg (determined by BCA method, BSA was used as a standard)

Formulation:

The antibody is lyophilized in 0.05 M phosphate buffer, 0.1 M NaCl, pH 7.2. ****AZIDE FREE****.

Reconstitution:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely. Slight turbidity may occur after reconstitution, which does not affect activity of the antibody. In this case clarify the solution by centrifugation.

Shipping:

At ambient temperature. Upon receipt, store the product at the temperature recommended below.

Storage/Stability:

The lyophilized antibody remains stable and fully active until the expiry date when stored at -20°C . Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles and store frozen at -80°C . Reconstituted antibody can be stored at 4°C for a limited period of time; it does not show decline in activity after one week at 4°C .

Quality Control:

Indirect ELISA – to determine titer of the antibody

SDS PAGE – to determine purity of the antibody

BCA - to determine quantity of the antibody

Applications:

Western blotting

Note:

This product is for research use only.